

Isaac Chemistry Skills

Developing mastery of essential pre-university physical chemistry

D.I. FOLLOWS
Head of Chemistry
Winchester College



Periphyseos Press
Cambridge, UK.

PERIPHYSEOS PRESS
Cambridge

Cavendish Laboratory
J. J. Thomson Avenue, Cambridge CB3 0HE, UK

Published in the United Kingdom by Periphyseos Press, Cambridge
www.periphyseos.org.uk

Developing mastery of essential pre-university physical chemistry
© D.I. Follows 2016
Creative Commons Licence

Developing mastery of essential pre-university physical chemistry is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Periphyseos Press.

First published 2016. First & second reprints, 2017.

Printed and bound in the UK by Short Run Press Limited, Exeter.

Typeset in L^AT_EX

A catalogue record for this publication is available from the British Library

ISBN 978-0-9572873-5-8 Paperback

Use this book in parallel with the electronic version at isaacchemistry.org. Marking of answers and compilation of results is free on Isaac Chemistry. Register as a student or as a teacher to gain full functionality and support.



used with kind permission of M. J. Rutter.
'The Periodic Table of the Elements' reproduced with kind permission of OCR

Contents

Notes for the Student and the Teacher	iv
Using Isaac Chemistry with this book; Significant figures	v
Mapping of Book Sections to Exam Specifications	vi
Table of Constants	viii
A Formulae & Equations	1
A1 Empirical formulae	1
A2 A_r & M_r and molecular formula	2
B Amount of Substance	3
B1 Standard form	3
B2 Unit conversions	4
B3 Gases	5
B4 Solids	6
B5 Solutions	6
B6 Reactions	7
B7 Titration	8
B8 Parts per million	11
C Gas Laws	12
C1 Ideal gases	12
C2 Density	15
C3 Non-ideal gases	16
D Atomic Structure	17
D1 Atomic structure	17
D2 Atomic orbitals	18

D3	Atomic and ionic radii and ionization energy	19
D4	Isotopes	21
E	Electronic Spectroscopy	23
F	Enthalpy Changes	25
F1	Calorimetry	25
F2	Bond enthalpies	28
F3	Formation and combustion enthalpies	29
F4	Born-Haber cycles	31
G	Entropy	35
G1	Absolute entropy	35
G2	Entropy changes	36
H	Free Energy	37
H1	Entropy change of the surroundings	37
H2	Free energy changes	39
H3	Free energy cycles	40
I	Equilibrium	43
I1	Equilibrium constant, K_p	43
I2	Equilibrium constant, K_c	45
I3	Solubility product	49
I4	Partition	51
I5	$RT \ln K$	53
J	Acids & Bases	55
J1	Brønsted-Lowry & Lewis	55
J2	pH & K_w	56
J3	K_a & pK_a	57
J4	K_b & pK_b	58
J5	Buffers	59
K	Redox	61
K1	Oxidation number	61
K2	Half-equations	62
K3	Balancing redox equations	63
K4	Disproportionation	65
L	Electrochemistry	66

L1	Electrode potential & cell potential	66
L2	Free energy & K_c	67
M	Rate Laws, Graphs & Half-life	68
M1	Rate laws	68
M2	Half-life	73
M3	The Arrhenius model	76
M4	Catalysis	79
M5	Michaelis-Menten kinetics	81

NOTES FOR THE STUDENT AND THE TEACHER

- “Developing mastery of essential pre-university physical chemistry” is associated with the MOOC: isaacchemistry.org
- The problems in this book are designed, graded and arranged by sections for understanding and mastery of the concepts and problem solving required at A-level chemistry and for equivalent exams.
- A mapping of each section of the book on to the relevant parts of all major UK chemistry exam specification is given on pages vi and vii. [With thanks to Dr David Paterson.] On-line, the mapping at isaacchemistry.org/pages/syll_map_chem allows 2-click setting of a chosen section as homework. See also below for teacher functionality.
- Each question can be answered on the above MOOC where there is immediate marking and some feedback. Practice to achieve *mastery* will give fluency and confidence in the essential skills required for strong A-level results and in the transition to university. Mastery level, that is achieving at least 80% of questions correct, is indicated in the marginal square boxes .
- Students should register (free) on Isaac. Go to isaacchemistry.org – click the “log in” tab and then “sign up” if not already registered. Physics registration on isaacphysics.org carries over for chemistry, and *vice versa*. A benefit of registration is that all activity is recorded for the student. Should questions be later set as homework, they will automatically be noted as done. All activity is confidential to the student, except parts that the student decides to share with a teacher (e.g. set home work).
- Teachers who register can have their Isaac accounts converted to teacher accounts (see at the foot of the user profile page). They can then set home work, and have it automatically marked with results returned to them: See “For Teachers” at isaacchemistry.org or go directly to isaacchemistry.org/teacher_features. Chemistry and physics functionalities are the same and are activated by registration on either the Physics or Chemistry sites.
- Visit isaacchemistry.org for physical chemistry problems in addition to these chemistry book problems (currently at “Further problem solving”). Be sure to select the level required, or otherwise get questions from all levels. These questions are all chemical thermodynamics for pre-AS, AS, A2, & the transition to university, with reaction kinetics soon to follow.
- See the physics companion book, also at www.isaacbooks.org/. It too is available for £1 per copy. Problems from the physics book can be answered on-line at isaacphysics.org/physics_skills_14.
- Maths problems to give the fluency required at chemistry and physics A-level, and equivalents, are available on isaacphysics.org.

USING ISAAC CHEMISTRY WITH THIS BOOK

Isaac Chemistry offers on-line versions of each sheet at isaacchemistry.org/book16 where a student can enter answers. This on-line tool will mark answers, giving immediate feedback to a student who, if registered on isaacchemistry.org, can have their progress stored and even retrieved for their CV! Teachers can set a sheet for class homework as the appropriate theme is being taught, and again for pre-exam revision. Isaac Chemistry can return the fully assembled and analysed marks to the teacher, if registered for this free service. Isaac Chemistry zealously follows the significant figures (sf) rules below and warns if your answer has a sf problem.

UNCERTAINTY AND SIGNIFICANT FIGURES

In chemistry, numbers represent values that have uncertainty and this is indicated by the number of significant figures in an answer.

Significant figures

When there is a decimal point (dp), all digits are significant, except leading (leftmost) zeros: 2.00 (3 sf); 0.020 (2 sf); 200.1 (4 sf); 200.010 (6 sf)

Numbers without a dp can have an *absolute accuracy*: 4 people; 3 electrons.

Some numbers can be ambiguous: 200 could be 1, 2 or 3 sf (see below). Assume such numbers have the same number of sf as other numbers in the question.

Combining quantities

Multiplying or dividing numbers gives a result with a number of sf equal to that of the number with the smallest number of sf:

$x = 2.31$, $y = 4.921$ gives $xy = 11.4$ (3 sf, the same as x).

An absolutely accurate number multiplied in does not influence the above.

Standard form

On-line, and sometimes in texts, one uses a letter 'x' in place of a times sign and ^ denotes "to the power of":

1800000 could be 1.80×10^6 (3 sf) and 0.0000155 is 1.55×10^{-5}
(standardly, 1.80×10^6 and 1.55×10^{-5})

The letter 'e' can denote "times 10 to the power of": 1.80e6 and 1.55e-5.

Significant figures in standard form

Standard form eliminates ambiguity: In $n.mnn \times 10^n$, the numbers before and after the decimal point are significant:

191 = 1.91×10^2 (3 sf); 191 is $190 = 1.9 \times 10^2$ (2 sf); 191 is $200 = 2 \times 10^2$ (1 sf).

Answers to questions

In this book and on-line, give the appropriate number of sf: For example, when the least accurate data in a question is given to 3 significant figures, then the answer should be given to three significant figures; see above. Too many sf are meaningless; giving too few discards information. Exam boards also require consistency in sf.

	OCR A (H032/H432)	OCR B (H033/H433)	AQA (7404/7405)	Edexcel (8CH0/9CH0)	Eduqas (A410QS)	CIE Pre-U (9791)	IB Chemistry (Higher)
A Formulae & Equations							
A1 Empirical formulae	2.1.3	EL(b)	3.1.2.4	Topic 5	C1.3	A4.1	1.2
A2 A_r & M_r and molecular formula	2.1.3	EL(a)	3.1.1.2/3.1.2.1	Topic 1	C1.3	A4.1	1.2
B Amount of Substance							
B1 Standard form	throughout	throughout	throughout	throughout	throughout	throughout	throughout
B2 Unit conversions	throughout	throughout	throughout	throughout	throughout	throughout	throughout
B3 Gases	2.1.3	DF(a)	3.1.2.2	Topic 5	C1.3	A1.1	1.2
B4 Solids	2.1.3	EL(b)	3.1.2.2	Topic 5	C1.3	A1.1	1.2
B5 Solutions	2.1.3	DF(a)	3.1.2.2	Topic 5	C1.3	A4.1	1.3
B6 Reactions	2.1.3	EL(b)	3.1.2.5	Topic 5	C1.3	A1.1	1.3
B7 Titration	2.1.4	EL(c)	3.1.12.5	Topic 5	C1.3	A4.1	1.3
B8 Parts per million	2.1.3	OZ(i)					1.3
C Gas Laws							
C1 Ideal gases	2.1.3	DF(a)	3.1.2.3	Topic 5	C1.3	B1.7	1.3
C2 Density							1.3
C3 Non-ideal gases							1.3
D Atomic Structure							
D1 Atomic structure	2.2.1	EL(f), DM(h)	3.1.1.3	Topic 1	C1.2	A1.2	2.2
D2 Atomic orbitals	2.2.1	EL(e)	3.1.1.3	Topic 1	C1.2	A1.2	2.2
D3 Atomic and ionic radii and ionization energy	3.1.1	EL(q)	3.1.1.3	Topic 1	C1.2	A1.2	3.2
D4 Isotopes	2.1.1	EL(h)	3.1.1.2	Topic 1	C1.3	A1.1	2.1, C.3, D.8
E Electronic Spectroscopy							
Electronic Spectroscopy		EL(w)	3.2.5.4	Topic 1	C1.2	A4.3	2.2
F Enthalpy Changes							
F1 Calorimetry	3.2.1	DF(f)	3.1.4.2	Topic 8	C2.2	A1.4	5.1
F2 Bond enthalpies	3.2.1	DF(e)	3.1.4.4	Topic 8	C2.2	A1.4	5.3
F3 Formation and combustion enthalpies	3.2.1	DF(g)	3.1.4.3	Topic 8	C2.2	A1.4	5.1
F4 Born-Haber cycles	5.2.1	(O(b))	3.1.8.1	Topic 13A	P14.1	A1.4	15.1
G Entropy							
G1 Absolute entropy	5.2.2	O(g)	3.1.8.2	Topic 13B	P14.2	B1.5	15.2
G2 Entropy changes	5.2.2	O(g)	3.1.8.2	Topic 13B	P14.2	B1.5	15.2

	OCR A (H032/H432)	OCR B (H033/H433)	AOA (7404/7405)	Edexcel (8CH0/9CH0)	Eduqas (A410QS)	CIE Pre-U (9791)	IB Chemistry (Higher)
H Free Energy							
H1 Entropy change of the surroundings	5.2.2	O(f)	3.1.8.2	Topic 13B		B1.5	15.2
H2 Free energy changes	5.2.2		3.1.8.2	Topic 13B	P14.2	B1.5	15.2
H3 Free energy cycles						B1.5	
I Equilibrium							
I1 Equilibrium constant, K_p	5.1.2		3.1.10	Topic 11	P15	B1.6	
I2 Equilibrium constant, K_c	3.2.3, 5.1.2	ES(p), Cl(h)	3.1.6.2	Topic 11	C2.1, P15.1	B1.6	7.1, 17.1
I3 Solubility product		O(h)				B1.6	A.10
I4 Partition							D.3
I5 $RT \ln K$				Topic 13B		B1.5	15.2, 17.1
J Acids & Bases							
J1 Brønsted-Lowry & Lewis	5.1.3	O(i)	3.1.12.1	Topic 12	P15.2	B1.6	8.1, 18.1
J2 pH & K_w	5.1.3	O(l)	3.1.12.2 / 3.1.12.3	Topic 12	P15.2	B1.6	8.3
J3 K_a & pK_a	5.1.3	O(l)	3.1.12.4	Topic 12	P15.2	B1.6	18.2
J4 K_b & pK_b						(B1.6)	18.2
J5 Buffers	5.1.3	O(m)	3.1.12.6	Topic 12	P15.2	B1.6	18.2
K Redox							
K1 Oxidation number	2.1.5	ES(e), DM(c)	3.1.7	Topic 3	C1.1, C1.6	B1.6	9.1
K2 Half-equations	5.2.3	ES(d), DM(c)	3.1.7	Topic 3	P11.1	B1.6	9.1, 19.1
K3 Balancing redox equations	5.2.3	DM(c)	3.1.7	Topic 3	P11.2	A2.1, B1.6	19.1, A.2
K4 Disproportionation	3.1.3, 5.3.1		3.1.7	Topic 4B	P12.1	A2.2	
L Electrochemistry							
L1 Electrode potential & cell potential	5.2.3	DM(f)	3.1.11.1	Topic 14	P11.1	B1.6	19.1
L2 Free energy & K_c	(5.2.3)	(DM(f))		(Topic 14)	(P11.1)	B1.6	19.1
M Rate Laws, Graphs & Half-life							
M1 Rate laws	5.1.1	Cl(a)	3.1.9.1	Topic 16	P13	B1.7	16.1
M2 Half-life	5.1.1	Cl(b)		Topic 16		B1.7	
M3 The Arrhenius model	5.1.1	Cl(d)	3.1.9.1	Topic 16	P13	B1.7	16.2
M4 Catalysis	various incl. 3.2.2, 3.2.3, 5.1.2	various in DF, OZ, Cl, CM	various incl. 3.1.5, .6, .10	Topic 9, 16	various incl. C2.3, P12.2, P13	A1.4, B1.7	various incl. 6.1, 7.1, 16.1, A.3
M5 Michaelis-Menten kinetics		(PL(f))				(B1.7)	B.7

Table of Constants

Quantity	Magnitude	Unit
Gas constant (R)	8.31	$\text{J mol}^{-1} \text{K}^{-1}$
Electron volt (eV)	1.60×10^{-19}	J
Avogadro's number (N_A)	6.02×10^{23}	–
Boltzmann's constant	1.38×10^{-23}	J K^{-1}
Speed of light (<i>in vacuo</i>)	3.00×10^8	m s^{-1}
Atomic mass unit (u)	1.66054×10^{-27}	kg
Planck's constant (h)	6.63×10^{-34}	J s
Charge on electron	1.60×10^{-19}	C
0 °C	273.15	K
Specific heat capacity of water	4180	$\text{J kg}^{-1} \text{K}^{-1}$
Density of water at RTP	1.00	g cm^{-3}
Faraday constant	96485	C mol^{-1}

Chapter A

Formulae & Equations

The boxed fraction shows how many questions need to be answered correctly to achieve mastery.

10/12

A1 Empirical formulae

A1.1 Find the empirical formulae for the ten compounds (a)–(j) from the data given. No compound contains more than 15 atoms in total in its formula. Show all your working in neat, clearly-presented answers. All compositions are by mass.

- a) 35.0% Nitrogen, 5.0% Hydrogen, 60.0% Oxygen
- b) 90.7% Lead, 9.3% Oxygen
- c) 26.6% Potassium, 35.3% Chromium, 38.1% Oxygen
- d) 40.3% Potassium, 26.8% Chromium, 32.9% Oxygen
- e) 29.4% Vanadium, 9.2% Oxygen, 61.4% Chlorine
- f) 81.8% Carbon, 18.2% Hydrogen
- g) 38.7% Carbon, 9.7% Hydrogen, 51.6% Oxygen
- h) 77.4% Carbon, 7.5% Hydrogen, 15.1% Nitrogen
- i) 25.9% Nitrogen, 74.1% Oxygen
- j) 29.7% Carbon, 5.8% Hydrogen, 26.5% Sulphur, 11.6% Nitrogen, 26.4% Oxygen

Element	Atomic mass	Element	Atomic mass
Hydrogen	1.0	Chlorine	35.5
Carbon	12.0	Potassium	39.1
Nitrogen	14.0	Vanadium	50.9
Oxygen	16.0	Chromium	52.0
Sulphur	32.1	Lead	207.2

A1.2 Complete combustion of 6.4 g of compound K produced 8.8 g of carbon dioxide and 7.2 g of water. Calculate the empirical formula of K.

- A1.3 Complete combustion of 1.8 g of compound L produced 2.64 g of carbon dioxide, 1.08 g of water and 1.92 g of sulfur dioxide. Calculate the empirical formula of L.

8/10

A2 A_r & M_r and molecular formula

Assume that the mass of an isotope in amu to 3 s.f. is equal to its mass number.

- A2.1 Which isotope is used as the standard against which relative atomic masses are calculated?
- A2.2 Fluorine only occurs naturally in one isotope, ^{19}F , and has a relative atomic mass of 19.0 amu. Calculate the mass of a fluorine atom in kg.
- A2.3 Magnesium has the following natural isotopes: ^{24}Mg 78.6%; ^{25}Mg 10.1%; ^{26}Mg 11.3%. Calculate the relative atomic mass of magnesium.
- A2.4 The relative atomic mass of boron is 10.8 amu. Boron exists in two isotopes, ^{10}B and ^{11}B . Calculate the percentage abundance of ^{10}B .
- A2.5 Complete the table below:

ELEMENT	A_r	Isotope 1	Isotope 2	Isotope 3	Isotope 4
Bromine	(a)	^{79}Br 50.5%	^{81}Br 49.5%	n/a	n/a
Silver	107.9	(b) ^{107}Ag ?%	(c) ^{109}Ag ?%	n/a	n/a
Cerium	140.2	^{136}Ce 0.2%	^{138}Ce 0.2%	^{140}Ce 88.5%	(d) ? ^{142}Ce 11.1%

- A2.6 The relative molecular mass of compound M is 135 amu. M contains 3.7% hydrogen, 44.4% carbon and 51.9% nitrogen by mass. Find the molecular formula of M.
- A2.7 Complete combustion of compound N occurs in a stoichiometric ratio of 1 : 6 with oxygen gas. Complete combustion of 4.2 g of compound N produces 13.2 g of carbon dioxide and 5.4 g of water. Find the molecular formula of N.

Chapter B

Amount of Substance

B1 Standard form

12/14

B1.1 Complete the following calculations, giving the answers in standard form to the appropriate number of significant figures. For guidance on this and on how to enter your answers on isaacchemistry.org, consult page ii of this book.

a) 120×70

b) $5600 + 800 + 12 + 1100 + 320$

c) $\frac{95.0}{19000}$

d) $12000 + 84000 + (3.00 \times 10^3) + 29000$

e) $(4.0 \times 10^2) \times 100 \times 300$

f) $\frac{1.6 \times 10^{-8}}{6.4 \times 10^{-3}}$

g) $\frac{3.00 \times 10^8}{5.2 \times 10^{-7}}$

h) $2.12 \times 10^{12} \times 5.4 \times 10^6$

i) $1.4 \times 10^{-10} \times 1.4 \times 10^{-10} \times 2.2 \times 10^{-10}$

j) $1.6 \times 10^{-19} \times 6.0 \times 10^{23}$

k) $\frac{1.3 \times 10^{17}}{3.0 \times 10^8}$

l) $(1.4 \times 10^{-6})^3$

m) $\sqrt{2.5 \times 10^{14}}$

n) $\frac{2.0 \times 10^4 \times 1.2 \times 10^4}{(3.2 \times 10^6)^2}$

o) $\frac{1.1 \times 10^{-5} \times (-2) \times 3}{(9.6 \times 10^{-11} + 1.2 \times 10^{-10})}$

B2 Unit conversions

Use standard form where answers are outside the range 0.01 to 1000 units.

B2.1 Convert the following volumes into dm^3

- | | |
|-------------------------|-------------------------|
| a) 0.86 m^3 | f) 70 cl |
| b) 200 cm^3 | g) 1.6 mm^3 |
| c) 45 ml | h) 1100 cc |
| d) 120 m^3 | i) 2.2 km^3 |
| e) 0.064 nm^3 | j) 42.5 \AA^3 |

B2.2 Converts the following masses into g:

- | | |
|-------------|---------------------|
| a) 16.0 kg | d) 12 tonne |
| b) 120 mg | e) $54 \mu\text{g}$ |
| c) 0.004 kg | |

B2.3 Convert the following into standard SI units:

- | | |
|---------------------------|-----------------------------|
| a) 68 km h^{-1} | f) 5.0 h |
| b) 500 g | g) 740 nm |
| c) 24 dm^3 | h) 72 mN cm^{-1} |
| d) 20 mbar | i) 1014 mbar |
| e) -77°C | j) 13.8 g cm^{-3} |

B2.4 Give the results of the following calculations in standard SI units:

- Density = $250 \text{ g} / 400 \text{ cm}^3$
- Speed = $96 \text{ km} / 80 \text{ min}$
- Concentration = $2.50 \text{ mmol} / 40.0 \text{ cm}^3$ (use mol dm^{-3})
- Momentum = $4.0 \times 10^{-23} \text{ g} \times 900 \text{ m s}^{-1}$
- Pressure = $590 \text{ fN} / 10 \text{ nm}^2$
- Volume = $240 \text{ pm} \times 240 \text{ pm} \times 320 \text{ pm}$
- Amount = $2.0 \mu\text{mol dm}^{-3} \times 75 \mu\text{m}^3$
- Energy = $3.2 \times 10^{-19} \text{ C} \times 2.4 \text{ kV}$

B3 Gases

RTP = room temperature and pressure.

Any gas occupies 24 dm^3 per mole at RTP.

Avogadro's number, $N_A = 6.02 \times 10^{23}$.

- B3.1 Calculate the volume occupied by:
- 4.0 moles of gas at RTP
 - 0.030 moles of gas at RTP
 - 5.0×10^{18} atoms of helium gas at RTP
 - 1.20×10^{24} molecules of ozone at RTP
 - 8.0 g of O_2 at RTP
 - 1.1 kg of carbon dioxide at RTP
- B3.2 Calculate the amount of gas (at RTP) in:
- 4.8 dm^3
 - 12 m^3
 - 400 cm^3
 - 18 ml
- B3.3 Calculate the number of molecules of gas (at RTP) in:
- 36 dm^3
 - 300 cm^3
- B3.4 Calculate the number of atoms (at RTP) in:
- 60 cm^3 of argon
 - 1.2 dm^3 of N_2
 - 8.0 m^3 of carbon dioxide
 - 420 cm^3 of ethene
- B3.5 Calculate the the mass of:
- 1.0 m^3 of neon at RTP
 - 20 cm^3 of $(\text{CH}_3)_2\text{O}$ at RTP
 - 420 cm^3 of ammonia at RTP

18/22

B4 Solids

B4.1 Find the molar masses in amu of the following compounds:

- | | |
|-----------------------------|--|
| a) CaCO_3 | f) FeSO_4 |
| b) Na_2CO_3 | g) KMnO_4 |
| c) NaOH | h) $\text{Fe}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ [1] |
| d) HCl | i) Calcium hydroxide |
| e) H_2SO_4 | j) Butane |

B4.2 Calculate the mass of:

- | | |
|---|--|
| a) 0.25 moles of $\text{H}_2\text{O}_2(\text{l})$ | d) 20.0 moles of $\text{Sr}(\text{s})$ |
| b) 6.0 moles of $\text{C}_2\text{H}_6(\text{g})$ | e) 1.20 moles of aluminium oxide |
| c) 0.40 moles of $\text{H}_2\text{O}(\text{l})$ | f) 7.4 moles of ammonium sulfate |

B4.3 Calculate the amount of:

- | | |
|---|--------------------------------------|
| a) 1.001 g of $\text{CaCO}_3(\text{s})$ | d) 2.006 kg of $\text{Hg}(\text{l})$ |
| b) 197 kg of $\text{Au}(\text{s})$ | e) 11.1 g of lithium carbonate |
| c) 1.4 g of $\text{CO}(\text{g})$ | f) 10.0 mg of lead(II) iodide |

10/12

B5 Solutions

B5.1 Calculate the concentration in mol dm^{-3} of the following solutions:

- 0.40 g NaOH in 100 ml water
- 7.3 g HCl in 1000 ml water
- 2.5 g H_2SO_4 in 50 ml water
- 15 g FeSO_4 in 500 ml water
- 0.16 g KMnO_4 in 200 ml water

B5.2 Calculate the mass of solute in each of the following:

- 500 ml of $0.010 \text{ mol dm}^{-3}$ NaOH
- 150 ml of 4.0 mol dm^{-3} HCl

[1]The 5 means that the formula includes 5 of what follows, i.e. water, so total mass is for $\text{Fe}_2\text{O}_3 + 5 \times \text{H}_2\text{O}$

- c) 1.00 ml of $10.0 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$
- d) 25.0 ml of $0.50 \text{ mol dm}^{-3} \text{ FeSO}_4$
- e) 21.8 ml of $0.0050 \text{ mol dm}^{-3} \text{ KMnO}_4$
- f) 2.0 dm^3 of $0.10 \text{ mol dm}^{-3} \text{ NaCl}$
- g) 100 ml of limewater with a concentration of $0.00020 \text{ mol dm}^{-3}$

B6 Reactions

17/21

- B6.1 Calculate the amount of oxygen needed, and amount of carbon dioxide produced, in each of the following cases:
- a) $\text{C}_3\text{H}_8 + 5 \text{O}_2 \longrightarrow 3 \text{CO}_2 + 4 \text{H}_2\text{O}$, using 1.0 mole of C_3H_8
 - b) $\text{C}_2\text{H}_6\text{O} + 3 \text{O}_2 \longrightarrow 2 \text{CO}_2 + 3 \text{H}_2\text{O}$, using 0.2 moles of $\text{C}_2\text{H}_6\text{O}$
 - c) $2 \text{CO} + \text{O}_2 \longrightarrow 2 \text{CO}_2$, using 4.0 moles of CO
 - d) $\text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{O}_2 \longrightarrow 6 \text{CO}_2 + 6 \text{H}_2\text{O}$, using 0.040 moles of $\text{C}_6\text{H}_{12}\text{O}_6$
 - e) $\text{C}_2\text{H}_4\text{O}_2 + 2 \text{O}_2 \longrightarrow 2 \text{CO}_2 + 2 \text{H}_2\text{O}$, using 0.10 moles of $\text{C}_2\text{H}_4\text{O}_2$
- B6.2 Calculate the amount of water produced by complete combustion of the following in oxygen (you will need to write a balanced equation each time):
- a) 1.0 mole of pentane, C_5H_{12}
 - b) 2.5 moles of heptane, C_7H_{16}
 - c) 200 moles of hydrogen, H_2
 - d) 4.0 moles of butane
 - e) 0.0030 moles of methane
- B6.3 Write the equation for each reaction and hence calculate the amount of acid required for complete reaction in each of the following cases:
- a) 0.10 mol NaOH reacting with H_2SO_4
 - b) HCl reacting with 20 g of CaCO_3
 - c) 24 g CuO reacting with HNO_3
 - d) 5.6 g Fe reacting with HCl
 - e) 14.8 g of calcium hydroxide reacting with H_2SO_4

f) 10 g of magnesium oxide reacting with nitric acid

B6.4 Calculate the volume of $0.50 \text{ mol dm}^{-3} \text{ H}_2\text{SO}_4$ required to neutralize each of the following:

- a) 25.0 cm^3 of $1.0 \text{ mol dm}^{-3} \text{ NaOH}$
- b) 3.0 g CaCO_3
- c) 1.25 g ZnCO_3
- d) 4.03 kg MgO
- e) 100 cm^3 of $0.2 \text{ mol dm}^{-3} \text{ NH}_3(\text{aq})$

20/24

B7 Titration

B7.1 Nitric acid of unknown concentration was added to a burette. 25.00 cm^3 of potassium hydroxide solution at a concentration of $0.100 \text{ mol dm}^{-3}$ was transferred to a 250 cm^3 conical flask using a volumetric pipette. A few drops of methyl orange indicator were added to the flask.

- a) Give the colour of the indicator in the alkaline solution in the flask.

The nitric acid was added to the flask a little at a time until the resulting solution went pink. The whole process was repeated until concordant titres (within 0.10 cm^3) were obtained.


- b) The concentration of nitric acid was found to be $0.092 \text{ mol dm}^{-3}$. Calculate the titre obtained, in cm^3 .

B7.2 In an analysis, 2.50 g of an unknown carbonate were dissolved in 100 cm^3 of 1.00 mol dm^{-3} hydrochloric acid (an excess). The resulting solution was made up to 250 cm^3 in a volumetric flask. 25.00 cm^3 aliquots of this solution were titrated against $0.250 \text{ mol dm}^{-3}$ sodium hydroxide. Some of the results are shown below. Fill in the gaps in the table overleaf, and then calculate the quantities below to identify the cation.

TITRATION	Initial burette reading / cm ³	Final burette reading / cm ³	Titre / cm ³
ROUGH	0.60	25.10	(a)
1	0.15	(b)	24.10
2	(c)	25.25	24.45
3	1.35	25.45	(d)

- e) Calculate the average concordant titre.
- f) Calculate the amount of sodium hydroxide in that volume.
- g) Calculate the amount of hydrochloric acid in each aliquot.
- h) Calculate the initial amount of hydrochloric acid added to the carbonate.
- i) Calculate the amount of hydrochloric acid remaining after reaction.
- j) Calculate the amount of hydrochloric acid used in reaction with the carbonate.
- k) Calculate the amount of carbonate in 2.50 g.
- l) Calculate the molar mass of the carbonate.
- m) Identify the cation in the carbonate.

B7.3 All of the ozone in 5.00 m³ of air was reacted with 250 cm³ of potassium iodide solution:



The liberated iodine was titrated against a standard solution of sodium thiosulfate with a concentration of 0.0400 mol dm⁻³. 25.0 cm³ of the iodine solution was used in each titration. The results of the titration are shown in the table overleaf. Fill in the remaining titres, and then answer the questions which follow.

TITRATION	Initial burette reading / cm ³	Final burette reading / cm ³	Titre / cm ³
ROUGH	0.10	25.40	25.30
1	0.80	26.10	(a)
2	1.20	26.20	(b)
3	1.00	25.90	(c)

- d) Calculate the concentration of the iodine solution.
- e) Calculate the amount of ozone in the 5.00 m³ of air.
- f) Name the piece of apparatus that should be used to transfer the iodine solution into a conical flask, ready for titration.
- g) Name a suitable indicator for this titration, and give its colour change at the end point.

B7.4 Three students each prepare a standard solution by dissolving 10.6 g of solid from different bottles labelled 'sodium carbonate' in exactly 1 dm³ of water. They use this standard solution in a titration to determine the exact concentration of a solution of sulphuric acid at approximately 0.1 mol dm⁻³. They each use a pipette to measure out exactly 25.00 cm³ of the standard solution into a conical flask; they use the same indicator and carry out their titrations with great care and accuracy. The volumes of sulphuric acid solution that they each use are tabulated below. Only student A finds the correct concentration of the sulphuric acid. Student B is within 20% but student C is so far out that they know something is wrong. Student C asks for help and is reminded that some solids can contain water of crystallization. Student A uses anhydrous sodium carbonate, but what is x in the formula Na₂CO₃ · x H₂O(s) for students B and C?

	Student A	Student B	Student C
Volume	23.75 cm ³	20.20 cm ³	8.80 cm ³

- a) Calculate the exact concentration of the sulphuric acid.
- b) Find x for the two different cases of students B and C.

B8 Parts per million

- B8.1 Calculate the ppm by volume of:
- 20 cm³ of CO per 40 m³ of air
 - 0.10 ml of alcohol per 100 ml of blood
 - 5.0 cm³ of O₃ per 20 m³ of air
 - 0.0040 cm³ of C₂H₄ per 1 dm³ of air
- B8.2 Calculate the ppm by mass of:
- 10 mg of Hg per tonne of water
 - 0.020 g of Mg per kg of CaCO₃
 - 50 mg of iron per kg of blood
 - 4.0×10^{-4} moles of arsenic per 1 kg of iron ore
- B8.3 Calculate the ppm by number of particles of:
- 23 mg of sodium in 2 kg of mercury
 - 60 μmol of albumen in 36 cm³ of water
 - 12 μg of magnesium hydrogen phosphate in 90 μl of water
 - 84 μg of carbon monoxide in 12 dm³ of air
- B8.4 Convert the following concentrations from parts per million (ppm) by mass to mol kg⁻¹.
- 2500 ppm CaCO₃
 - 32.0 ppm NH₃
 - 120 ppm H₂O₂
 - 0.25 ppm Hg
 - 6.0 ppm CH₃CH₂CH₂COOH

The Periodic Table of the Elements

(1) 1

(2)

(3)

(4)

(5)

(6)

(7)

(8)

Key
atomic number
Symbol
name
relative atomic mass

1	H	hydrogen	1.0
---	----------	----------	-----

2			
---	--	--	--

3	Li	lithium	6.9
4	Be	beryllium	9.0

11	Na	sodium	23.0
12	Mg	magnesium	24.3

19	K	potassium	39.1
20	Ca	calcium	40.1

37	Rb	rubidium	85.5
38	Sr	strontium	87.6

55	Cs	caesium	132.9
56	Ba	barium	137.3

87	Fr	francium	88
88	Ra	radium	88

21	Sc	scandium	45.0
22	Ti	titanium	47.9

39	Y	yttrium	88.9
40	Zr	zirconium	91.2

41	Nb	niobium	92.9
42	Mo	molybdenum	95.9

43	Tc	technetium	
44	Ru	ruthenium	101.1

45	Rh	rhodium	102.9
46	Pd	palladium	106.4

47	Ag	silver	107.9
48	Cd	cadmium	112.4

73	Ta	tantalum	180.9
74	W	tungsten	183.8

75	Re	rhenium	186.2
76	Os	osmium	190.2

77	Ir	iridium	192.2
78	Pt	platinum	195.1

79	Au	gold	197.0
80	Hg	mercury	200.6

81	Tl	thallium	204.4
82	Pb	lead	207.2

13	B	boron	10.8
14	C	carbon	12.0

13	Al	aluminium	27.0
14	Si	silicon	28.1

15	P	phosphorus	31.0
16	S	sulfur	32.1

31	Ga	gallium	69.7
32	Ge	germanium	72.6

33	As	arsenic	74.9
34	Se	selenium	79.0

51	Sb	antimony	121.8
52	Te	tellurium	127.6

83	Bi	bismuth	208.98
84	Po	polonium	

85	At	astatine	
86	Rn	radon	

101	Md	mendelevium	
102	No	nobelium	

103	Lr	lawrencium	
-----	-----------	------------	--

109	Mt	meitnerium	
110	Ds	darmstadtium	

111	Rg	roentgenium	
112	Cn	copernicium	

114	Fl	flerovium	
116	Lv	livermorium	

117			
-----	--	--	--

17	F	fluorine	19.0
18	Ne	neon	20.2

35	Br	bromine	79.9
36	Kr	krypton	83.8

53	I	iodine	126.9
54	Xe	xenon	131.3

85	At	astatine	
86	Rn	radon	

9	F	fluorine	19.0
10	Ne	neon	20.2

17	Cl	chlorine	35.5
18	Ar	argon	39.9

2	He	helium	4.0
---	-----------	--------	-----

57	La	lanthanum	138.9
89	Ac	actinium	227.0

59	Pr	praseodymium	140.9
91	Pa	protactinium	231.0

61	Pm	promethium	
93	Np	neptunium	237.0

62	Sm	samarium	150.4
94	Pu	plutonium	244.0

63	Eu	euroium	152.0
95	Am	americium	243.0

64	Gd	gadolinium	157.2
96	Cm	curium	247.0

65	Tb	terbium	158.9
97	Bk	berkelium	247.0

66	Dy	dysprosium	162.5
99	Cf	californium	251.0

67	Ho	holmium	164.9
101	Md	mendelevium	288.1

68	Er	erbium	167.3
102	No	nobelium	289.1

69	Tm	thulium	168.9
103	Lr	lawrencium	260.1

70	Yb	ytterbium	173.0
104	Lr	lawrencium	261.1

71	Lu	lutetium	175.0
----	-----------	----------	-------